

Interaction Effects on the Conductance in One-Dimensional Systems – Short-Range Interaction^{*}

Arisato KAWABATA[†]

Department of Physics, Gakushuin University,
1-5-1 Mejiro, Toshima-ku, Tokyo 171

February 7, 2008

Abstract

We investigate the effect of electron-electron interactions on the conductance of quasi one-dimensional systems without potential scattering. For a finite temperature or system length, the short-range interaction is not renormalized to 0, and it gives rise to a finite correction to the conductance if we calculate it using Kubo formula. We show that this correction can be absorbed into the renormalization of the chemical potential and that the properly defined conductance to be observed in the experiments is equal to that of non-interacting electrons.

keyword: Tomonaga-Luttinger liquid, one-dimensional system quantum wire, Coulomb interaction

1 Introduction

Electron transport in one-dimensional interacting electron system is not yet fully understood in spite of its simplicity. Even in the absence of the potential scattering, until recently it has been believed that the conductance is renormalized by the long-range interaction like $2Ke^2/h$, where $K < 1$ for repulsive interaction.[1] The experiment by Tarucha et al. ,[2] however, indicates that the renormalization is absent, and it has been shown by the present author that the renormalization is the result of the incorrect definition of the conductance.[3]

He showed that the renormalization can be derived with the use of the self-consistent field method: When a electric potential is applied on the system,

^{*}submitted to J. Phys. Soc. Jpn.

[†]E-mail: arisato.kawabata@gakushuin.ac.jp

the electrons are driven by the self-consistent potential which is reduced because of the screening, while the current is defined as the ratio of the current to the difference of the externally applied potential between the ends of the system. This wrong definition of the conductance leads to the apparent reduction of the conductance. In fact, as has been pointed out by Izuyama,[4] the conductance should be defined as the ratio of the current to the difference of the self-consistent potential between the ends of the system, because the self-consistent potential which contains the contributions of the polarization charge is the electric potential observed in the experiments. Within the self-consistent field theory, which is equivalent to the exact theory in the present case, the response to the self-consistent potential is that of non-interacting electron, and hence the correctly defined conductance is not renormalized.

Here it should be noted that the above arguments are correct only when the long-range parts of the interaction potential are taken into account. It has been argued that the short-range parts are renormalized to 0,[1] but it is the case only for 0K and infinitely long systems. In fact, in order to determine whether the renormalization exists, the experiments have to be done at moderate temperatures to avoid the effects of the scattering by impurities or boundary irregularity.

Thus it is important to investigate the effects of the short-range part of the interaction on the conductance, and it is the purpose of this paper.

2 Correction to the Conductance

We consider a one-dimensional interacting electron system described by the Hamiltonian

$$\mathcal{H} = \sum_{k,\sigma} \frac{\hbar^2 k^2}{2m} a_{k,\sigma}^\dagger a_{k,\sigma} + \frac{1}{L} \sum_{p,k,q} V(q) a_{p+q,\uparrow}^\dagger a_{k-q,\downarrow}^\dagger a_{k,\downarrow} a_{p,\uparrow}. \quad (1)$$

Suppose we apply an electric field $E(x)e^{\delta t}$ ($\delta \rightarrow +0$) along the system, then the current is given by[5]

$$I(x) = \int \sigma(x, x') E(x') dx', \quad (2)$$

$$\sigma(x, x') = \lim_{\delta \rightarrow +0} \int_0^\beta d\lambda \int_0^\infty dt \langle j(x', -i\hbar\lambda) j(x, t) \rangle e^{\delta t}, \quad (3)$$

where $j(x, t)$ is the Heisenberg representation of the current operator. It is easy to see that eq. (3) can be rewritten in the form

$$\sigma(x, x') = \lim_{\delta \rightarrow +0} \frac{1}{\delta} \{ G(x, x', \delta) - G(x, x', 0) \}, \quad (4)$$

$$G(x, x', \omega_n) = - \int_0^{\hbar\beta} \langle \bar{j}(x, \tau) j(x') \rangle e^{i\omega_n \tau / \hbar} d\tau, \quad (5)$$

with

$$\bar{j}(x, \tau) = e^{\mathcal{H}\tau/\hbar} j(x) e^{-\mathcal{H}\tau/\hbar}, \quad (6)$$

where $\omega_n = 2\pi k_B T n$, n being an integer, and $G(x, x', \delta)$ is obtained by analytic continuation.

We let $G(q, \omega_n)$ be the Fourier transform of $G(x, x', \omega_n)$;

$$G(x, x', \omega_n) = \int \frac{dq}{2\pi} G(q, \omega_n) e^{iq(x-x')}. \quad (7)$$

In the lowest order in the short-range interaction, there are two kinds of corrections to $G(q, \omega_n)$, namely, vertex correction (Fig.1(a)), and self-energy correction (Fig. 1(b)).




Figure 1: Vertex correction (a), and self-energy correction (b) to $G(q, \omega_n)$. The solid and the dotted lines indicate the electron Green's functions and the interactions, respectively.

The vertex correction is given by

$$\begin{aligned} \Delta G_v(q, \omega_n) = & -2\hbar(k_B T)^2 \sum_{\varepsilon_n, \varepsilon_m} \int \frac{dk}{2\pi} \int \frac{dp}{2\pi} V(p-k) \left(\frac{e\hbar}{2m} \right)^2 \\ & \times (2k+q)(2p+q) \mathcal{G}(k, \varepsilon_n) G(k+q, \varepsilon_n + \omega_n) \mathcal{G}(p, \varepsilon_m) \mathcal{G}(p+q, \varepsilon_m + \omega_n), \end{aligned} \quad (8)$$

where $\mathcal{G}(k, \varepsilon_n)$ is the one-electron Green's function;

$$\mathcal{G}(k, \varepsilon_n) = \frac{1}{i\varepsilon_n - \xi_k}, \quad (9)$$

with $\varepsilon_n = (2n+1)k_B T$, $\xi_k = \hbar^2(k^2 - k_F^2)/2m$ and k_F being the Fermi wave number. Using the standard technique of the thermal Green's function method,[6] we find that

$$\begin{aligned} \Delta G_v(q, \omega_n) = & -2\hbar \left(\frac{e\hbar}{2m} \right)^2 \int \frac{dk}{2\pi} \int \frac{dp}{2\pi} V(p-k) \\ & \times (2k+q)(2p+q) \frac{f(\xi_{k+q}) - f(\xi_k)}{\xi_{k+q} - \xi_k - i\omega_n} \frac{f(\xi_{p+q}) - f(\xi_p)}{\xi_{p+q} - \xi_p - i\omega_n}, \end{aligned} \quad (10)$$

where $f(\xi)$ is the Fermi distribution function.

We assume that the spatial change of the electric field $E(x)$ is very slowly, and we calculate $\Delta G_v(q, \omega_n)$ for $q \ll k_F$. Then, for $k_B T \ll \varepsilon_F$, ε_F being the Fermi energy, in the right hand side of eq. (10) only the wave number $k, p \approx \pm k_F$ contribute to the integrals. Since we are interested in the correction due to the short range part of $V(q)$, we consider only the contributions from $|k - p| \approx 2k_F$. Then we easily find that

$$\Delta G_v(q, \omega_n) = \left(\frac{ev_F}{\pi} \right)^2 V(2k_F) \frac{\hbar q^2}{(\hbar v_F q)^2 + \omega_n^2}, \quad (11)$$

and from eqs. (4) and (7) that the vertex correction to the conductivity is given by

$$\begin{aligned} \Delta \sigma_v(x, x') &= - \lim_{\delta \rightarrow +0} \frac{e^2}{\pi^2 \hbar} V(2k_F) \int \frac{\delta e^{iq(x-x')}}{(\hbar v_F q)^2 + \delta^2} \frac{dq}{2\pi} \\ &= - \frac{2e^2}{h} \frac{V(2k_F)}{h v_F}. \end{aligned} \quad (12)$$

Next we calculate the self-energy correction. The correction of Fig.1(b) is given by

$$\begin{aligned} \Delta G_{s1}(q, \omega_n) &= 2\hbar k_B T \sum_{\varepsilon_n} \int \left(\frac{e\hbar}{2m} \right)^2 (2k + q)^2 \\ &\quad \times \mathcal{G}^2(k, \varepsilon_n) \Sigma(k) \mathcal{G}(k + q, \varepsilon_n + \omega_n) \frac{dk}{2\pi}, \end{aligned} \quad (13)$$

where $\Sigma(k)$ is the self-energy part, which is independent of the energy :

$$\begin{aligned} \Sigma(k) &= - k_B T \sum_{\varepsilon_m} \int V(k - p) \mathcal{G}(p, \varepsilon_m) \frac{dp}{2\pi} \\ &= - \int V(k - p) f(\xi_p) \frac{dp}{2\pi}. \end{aligned} \quad (14)$$

Together with the contribution of the Feynman graph with the self-energy correction to the other one-electron Green's function in Fig.1(b), we find that the self-energy correction to $G(q, \omega_n)$ is given by

$$\begin{aligned} \Delta G_s(q, \omega_n) &= -2\hbar \left(\frac{e\hbar}{2m} \right)^2 \int (2k + q)^2 \\ &\quad \times \left[\frac{\{f(\xi_{k+q}) - f(\xi_k)\} \{\Sigma(k + q) - \Sigma(k)\}}{(\xi_{k+q} - \xi_k - i\omega_n)^2} + \frac{f'(\xi_{k+q}) - f'(\xi_k)}{\xi_{k+q} - \xi_k - i\omega_n} \right] \frac{dk}{2\pi}, \end{aligned} \quad (15)$$

where the primes on $f(\xi)$ denote the derivative with respect to ξ .

The self-energy correction to the conductivity is obtained from eqs. (4) and (7) with $G(q, \omega_n)$ replaced by $\Delta G_s(q, \omega_n)$ in eq. (15). After some manipulations we find that

$$\Delta\sigma_s(x, x') = \int \frac{e^2 \hbar |k|}{m} [f'(\xi_k) \Sigma'(k) - f''(\xi_k) \Sigma(k)] \frac{dk}{2\pi}, \quad (16)$$

where the primes indicate the derivative with respect to ξ_k . It is easy to see that this equation can be written in the form

$$\Delta\sigma_s(x, x') = \frac{4e^2}{\hbar} \int_0^\infty f'(\xi_k) \frac{d\Sigma(\xi_k)}{dk} \frac{dk}{2\pi}. \quad (17)$$

The derivative of the self-energy part is obtained from eq. (14):

$$\frac{d\Sigma(k)}{dk} = - \int V(k-p) \frac{df(\xi_p)}{dp} \frac{dp}{2\pi}, \quad (18)$$

and for $k_B T \ll \varepsilon_F$, it follows that

$$\Delta\sigma_s(x, x') = \frac{2e^2}{h} \frac{2V(2k_F)}{hv_F}, \quad (19)$$

where we have neglected the contribution of the term with $V(q)$ for $q \ll 2k_F$.

Thus together with eq. (12) the correction to the conductivity to the first order in the short-range part of the interaction is given by

$$\Delta\sigma(x, x') = \frac{2e^2}{h} \frac{V(2k_F)}{hv_F}. \quad (20)$$

Since the right hand side is independent of x and x' , we find that the correction to the conductance is also given by the right hand side of the above equation.

3 Renormalization of the Chemical Potential

In this section we show that correction to the conductance obtained in the preceding section can be interpreted as the one due to the correction to the chemical potential difference at the ends of the sample.

The electric potential applied on the system causes the deviation $\Delta n(x)$ of the electron density from its unperturbed value, and it gives rise to the change in the local chemical potential.

We calculate $\Delta n(x)$ as the response to the electric potential $\Phi(x)e^{\delta t/\hbar}$ ($\delta \rightarrow +0$), with $E(x) = -d\Phi(x)/dx$, and we obtain

$$\Delta n(x) = \lim_{\delta \rightarrow 0} \int \Phi(q) R_0(q, \delta) e^{iqx} \frac{dq}{2\pi}, \quad (21)$$

where $\Phi(q)$ is the Fourier component of $\Phi(x)$ and $R_0(q, \delta)$ is the density response function:[3]

$$R_0(q, \delta) = \frac{2ev_F q^2}{\pi[(\hbar v_F q)^2 + \delta^2]}. \quad (22)$$

Then we obtain

$$\Delta n(x) = \frac{2e}{\pi \hbar v_F} \Phi(x). \quad (23)$$

Note that the density of states per spin is $1/\pi \hbar v_F$. Hence the local Fermi energy is given by

$$\Delta \varepsilon_F(x) = e\Phi(x). \quad (24)$$

Here the Fermi energy is measured from the bottom of the band.

On the other hand, the interaction correction to the chemical potential is given by

$$\Delta \mu = \Delta \Sigma(k_F) = - \int V(k_F - p) \Delta f(\xi_p) \frac{dp}{2\pi}, \quad (25)$$

because the change of the self-energy have to be compensated by that of the chemical potential in order to keep the electron density unchanged. Here we assume that the local relation between the Fermi energy and the chemical potential holds, i.e.,

$$\Delta \mu(x) = - \int V(k_F - p) \frac{\partial f(\xi_p)}{\partial \mu} \Delta \varepsilon_F(x) \frac{dp}{2\pi}, \quad (26)$$

and then, neglecting the contributions of $V(q)$ for $|q| \ll 2k_F$, it follows from eq. (24) that

$$\Delta \mu(x) = -e\Phi(x) \frac{V(2k_F)}{2\pi \hbar v_F}. \quad (27)$$

The increase of the current due to this correction to the chemical potential difference is given by

$$\begin{aligned} \Delta I &= \frac{2e^2}{h} \frac{\{\Delta \mu(-\infty) - \Delta \mu(\infty)\}}{-e} \\ &= \frac{2e^2}{h} \frac{V(2k_F)}{\hbar v_F} \{\Phi(-\infty) - \Phi(\infty)\}. \end{aligned} \quad (28)$$

Comparing it with eq. (20), we find that the correction to the conductance due to the short-range parts of the interaction potential can be explained in terms of the corrections to the chemical potential at the ends of the system. Therefore, the correction to the conductance can be absorbed into the renormalization of the chemical potential, i.e., the conductance is not renormalized if we define it as the ratio of the current to the difference of the renormalized chemical potential at the ends of the system.

4 Discussion

In the previous paper[3], the present author has pointed out that the renormalization of the conductance due to the long-range part of the interaction can be explained in terms of the renormalization of the electric potential. Since the renormalized potential is the one observed in the experiments, the conductance should be defined as the ratio of the current to the difference of the renormalized potential at the ends of the sample. The conductance defined in this way is not renormalized.

In this paper, using Kubo formula, we showed that the short-range part of the interaction gives a finite correction to the conductance and that it can be ascribed to the renormalization of the chemical potential. As in the above, the renormalized chemical potential is the one observed in the experiments, and the conductance should be defined in terms of the renormalized chemical potential. Then the conductance is not renormalized also in this case. Although the present arguments are restricted to the lowest order in the interaction potential, we can expect that it is the case to all order.

Recently, there are some calculations on the effects of the Umklapp scattering. In the Umklapp scattering, the total momentum of electrons is not conserved, and we can expect a finite correction to the conductance. From the point of view of the present theory, however, some contributions to the correction obtained using Kubo formula might be absorbed into the renormalization of the chemical potential. Such contributions are not to be observed in the experiments.

Acknowledgments

The author is grateful to Prof. T. Ando, Prof. A. Shimizu, and Prof. K. Kitahara for valuable discussions. This work is partly supported by Grant-in-Aid for Scientific Research on Priority Area "Mesoscopic Electronics: Physics and Technology" from the Ministry of Education, Science, Sports and Culture.

References

- [1] J. Solyom: Adv. Phys. **28** (1979) 201.
- [2] S. Tarucha, T. Honda and T. Saku: Solid State Commun. **94** (1995) 413.
- [3] A. Kawabata: J. Phys. Soc. Jpn. **65** (1996) 30.
- [4] T. Izuyama: Prog. Theor. Phys. **25** (1961) 964.
- [5] R. Kubo: J. Phys. Soc. Jpn. **12** (1957) 570.

- [6] A.A. Abrikosov, L.P. Gor'kov and I.E. Dzyaloshinskii: *Methods of Quantum Field Theory in Statistical Physics* (Pergamon Press, London, 1963, and Dover Publication Inc., New York, 1975)